

ИЗВЕСТИЯ

Гомельского государственного университета
имени Ф. Скорины

№ 6 (123)

Естественные науки

2020

Гомельский государственный университет
имени Ф. Скорины

ИЗВЕСТИЯ

Журнал зарегистрирован в Министерстве информа-
ции Республики Беларусь
(свидетельство о регистрации
№ 546 от 06.07.2009 года)

Журнал включен ВАК Республики Беларусь
в перечень научных изданий Республики Беларусь,
в которых публикуются результаты
диссертационных исследований
(приказы № 207 от 13.12.2005, № 9 от 15.01.2010,
№ 57 от 16.05.2013)

Журнал включен в библиографические базы данных
ВИНИТИ и Научную электронную библиотеку
eLIBRARY.RU

РЕДАКЦИОННАЯ КОЛЛЕГИЯ

Главный редактор С.А. ХАХОМОВ,
д-р. физ.-мат. наук, доцент
Зам. главн. редактора О.М. ДЕМИДЕНКО,
д-р тех. наук, профессор
Зам. главн. редактора М.В. СЕЛЬКИН,
д-р физ.-мат. наук, профессор

Члены редакционной коллегии:

Г.Г. Гончаренко, д-р биол. наук, проф.,
чл.-корр. НАН Беларуси
Ф.В. Кадол, д-р пед. наук, проф.
В.Н. Калмыков, д-р филос. наук, проф.
В.И. Коваль, д-р филол. наук, проф.
Г.Г. Лазько, д-р ист. наук, проф.
И.В. Семченко, д-р физ.-мат. наук, проф.
В.С. Смородин, д-р тех. наук, проф.
Б.В. Сорвилов, д-р экон. наук, проф.
В.М. Хомич, д-р юрид. наук, проф.
О.Г. Шляхтова, ответственный секретарь

Члены редакционной коллегии по естественным наукам:

В.Ф. Багинский, д-р биол. наук, проф., член-корр.
НАН Беларуси
А. Баллистер-Болинше (Испания), д-р, проф. математики
Ван Сяо Фэн (Китай), д-р тех. наук, проф.
Го Вэньбинь (Китай), д-р физ.-мат. наук, проф.
В.П. Кудин, д-р тех. наук, проф.
А.А. Махнев (Россия), д-р физ.-мат. наук, проф.,
член-корр. РАН
В.В. Можаровский, д-р тех. наук, проф.
А.Н. Сердюков, д-р физ.-мат. наук, проф., член-
корр. НАН Беларуси
А.Н. Скиба, д-р физ.-мат. наук, проф.
Шэн Рикун (Китай), д-р тех. наук, проф.
Р. Эстебан Ромеро (Испания), д-р, проф. математики

АДРЕС РЕДАКЦИИ:
246019, Беларусь, Гомель, ул. Советская, 104,
Телефоны: +375 (232) 51-03-21
E-mail: vesti@gsu.by
Интернет-адрес: <http://vesti.gsu.by>

Francisk Scorina Gomel State University

PROCEEDINGS

The Journal is registered in the Ministry of Information of
Republic of Belarus
(registration certificate
number 546 dated 06.07.2009)

The Journal is included in the Republic of Belarus
Higher Attestation Commission list of scientific publica-
tions of the Republic of Belarus, which publish the main
results for the degree of Doctor (Candidate) of Sciences
(order number 207 dated 13.12.2005, number 9 dated
15.01.2010, number 57 dated 16.05.2013)

The Journal is included in bibliographic databases of the
All-Russia Institute of Scientific and
Technical Information (VINITI), Scientific electronic
library eLIBRARY.RU

EDITORIAL BOARD

Editor-in-chief S.A. KHAKHOMOV,
Sc. D., Docent of Physics
Deputy editor-in-chief O.M. DEMIDENKO,
Sc. D., Professor
Deputy editor-in-chief M.V. SELKIN,
Sc. D., Professor

Members of editorial board:

G.G. Goncharenko, Sc. D., Professor, Corresponding
Member NASB
F. V. Kadol, Sc. D., Professor
V.N. Kalmykov, Sc. D., Professor
V.I. Koval, Sc. D., Professor
G.G. Lazko, Sc. D., Professor
I.V. Semchenko, Sc. D., Professor
V.S. Smorodin, Sc. D., Professor
B.V. Sorvirov, Sc. D., Professor
V.M. Homich, Sc. D., Professor
O.G. Shlyahova, executive secretary

Members of editorial board for the natural sciences

V.F. Baginsky, Sc. D., Professor, Corresponding
Member NASB
A. Ballister-Bolinshes (Spain), Sc. D., Professor
Van Siao Fen (China), Sc. D., Professor
Go Wenbin (China), Sc. D., Professor
V.P. Kudzin, Sc. D., Professor
A.A. Makhnev (Russia), Sc. D., Professor, Correspond-
ing Member RAS
V.V. Mozharovsky, Sc. D., Professor
A.N. Serdukov, Sc. D., Professor, Corresponding
Member NASB
A.N. Skiba, Sc. D., Professor
Shen Riku (China), Sc. D., Professor
R. Esteban Romero (Spain), Sc. D., Professor

EDITORIAL OFFICE ADDRESS:
246019, Belarus, Gomel, Sovetskaya Str., 104,
Tel: +375 (232) 51-03-21
E-mail: vesti@gsu.by
Site: <http://vesti.gsu.by>

ИЗВЕСТИЯ

Гомельского государственного университета
имени Ф. Скорины

НАУЧНЫЙ И ПРОИЗВОДСТВЕННО-ПРАКТИЧЕСКИЙ ЖУРНАЛ

Издается с 1999 г.
Выходит 6 раз в год

• 2020, № 6 (123) •

ЕСТЕСТВЕННЫЕ НАУКИ:

БИОЛОГИЯ • ТЕХНИКА (ИНФОРМАЦИОННЫЕ ТЕХНОЛОГИИ;
РАДИОТЕХНИКА И СВЯЗЬ; ЭЛЕКТРОНИКА) • МАТЕМАТИКА • ФИЗИКА

СОДЕРЖАНИЕ

Биология

Абрамова И.В. Статус птиц орнитокомплексов березовых лесов на различных стадиях сукцессий в юго-западной Беларуси	5
Автушко М.И. Об одном решении дилеммы Барбера	11
Багинский В.Ф., Демид Н.П. Повышение точности учета лесосечного фонда смешанных древостоев при лесоустройстве	18
Булко Н.И., Потапенко А.М., Митин Н.В., Бутьковец В.В., Козлов А.К., Толкачева Н.В., Машков И.А., Серенкова В.А. Состояние дочернобыльских сосновых лесов в 30-километровой зоне ЧАЭС	22
Дайнеко Н.М., Мироненко В.И., Мироненко В.В. Математическая модель влияния метеорологических условий на урожайность луговых экосистем поймы р. Сож	28
Дрозд К.С., Галиновский Н.Г. Сообщества щелкунов (Coleoptera, Elateridae) соснового леса в окрестностях скважин Южно-Осташковского нефтяного месторождения (Калинковичский район, Республика Беларусь)	34
Карпенко А.Ф. Состояние территории радиоактивного загрязнения и о мерах её возрождения	38
Концевая И.И. Влияние бета-лактамовых антибиотиков на органогенез в культуре тканей листовых эксплантов карельской березы	43
Лазарева М.С., Климович Л.К., Климов А.В. Особенности формирования дубравы кисличной	50
Логинов Д.Н. Видовая структура имаго кровососущих комаров (Diptera, Culicidae) на территории Брестской и Гомельской областей Республики Беларусь	56
Макаренко Т.В. Влияние природных сорбентов на накопление тяжелых металлов в донных отложениях водоемов	62
Островский А.М. Определитель синантропных видов мокриц (Isopoda: Oniscidea) юго-востока Беларуси	69
Сакович В.В., Жерносеков Д.Д. Молокосвертывающие ферменты различного происхождения: перспективы применения в сыроделии	75
Соколов А.С. Оценка ландшафтной репрезентативности системы особо охраняемых природных территорий Белорусской Возвышенной провинции	81
Углынец А.В., Гарбарук Д.К. Загрязнение ^{137}Cs и ^{90}Sr доминантных видов живого почвенного покрова в черноольшаниках зоны отчуждения Чернобыльской АЭС	85

Храмченкова О.М., Матвеевков М.В. <i>Цитотоксические и фотомодифицирующие свойства метанольных и ацетоновых экстрактов пяти видов лишайников</i>	92
ИНФОРМАТИКА	
Воруев А.В., Демиденко И.О., Колаиб С.М., Домасканов Д.В. <i>Построение рабочего места для тестирования виртуальной сетевой инфраструктуры</i>	99
Осипенко Н.Б., Мироненко В.М. <i>Автоматизация подбора персонифицированных предложений на примере задачи поиска фильма</i>	105
МАТЕМАТИКА	
Бородич Р.В., Селькин М.В. <i>О пересечении не \mathfrak{F}-подгрупп, выделяемых подгрупповым функтором</i>	111
Гальмак А.М., Селькин М.В. <i>Косые элементы в полиадических группах специального вида нечётной арности. II</i>	118
Малинковский Ю.В. <i>Критерий эргодичности для одного класса цепей Маркова с непрерывным временем</i>	125
Можей Н.П. <i>Эквивалентные связности нулевой кривизны на однородных пространствах с разрешимой группой преобразований</i>	131
Немилостивая В.А. <i>Стационарное распределение сетей с положительными и отрицательными заявками и двумя типами узлов</i>	138
Пилипчук Л.А., Пилипчук А.С., Полячок Е.Н. <i>Двухкритериальные задачи потокового программирования</i>	144
ФИЗИКА	
Леоненко Д.В. <i>Поперечный изгиб круговой сэндвич-пластины ступенчатой толщины</i> ...	151
Охрименко И.П., Кольчевский Н.Н., Петров П.В. <i>Модель геостационарного детектора гравитационных волн</i>	156
КРАТКИЕ СООБЩЕНИЯ	
Саварин А.А. <i>О находке белозубки малой (<i>Crocidura suaveolens</i>) на территории Могилевской области Беларуси</i>	163

PROCEEDINGS

of Francisk Scorina Gomel State University

SCIENTIFIC, PRODUCTION AND PRACTICAL JOURNAL

There are 6 times a year

Published since 1999

• 2020, № 6 (123) •

NATURAL SCIENCES:

BIOLOGY • TECHNIQUE (INFORMATION TECHNOLOGY;
RADIO ENGINEERING AND COMMUNICATIONS; ELECTRONICS)

• MATHEMATICS • PHYSICS

CONTENTS

BIOLOGY

Abramava I.V. <i>The status of birds of ornithocomplexes in birch forests at different stages of succession in southwestern Belarus</i>	5
Avtushko M.I. <i>One solution to Barber's dilemma</i>	11
Baginskij V.F., Demid N.P. <i>Improving the accuracy of accounting for the felling fund of mixed forest stands during forest management</i>	18
Bulko N.I., Potapenko A.M., Mitin N.V., But'kovec V.V., Kozlov A.K., Tolkacheva N.V., Mashkov I.A., Serenkova V.A. <i>The state of the pre-Chernobyl pine forests in the 30-kilometer zone of the Chernobyl nuclear power plant</i>	22
Dajneko N.M., Mironenko V.I., Mironenko V.V. <i>Mathematical model of the influence of meteorological conditions on the productivity of meadow ecosystems of the river Sozh</i>	28
Drozd K.S., Galinovskij N.G. <i>Communities of click-beetles (Coleoptera, Elateridae) of a pine forest in the vicinity of the wells of the Yuzhno-Ostashkovskoye oil field (Kalinkovichi district, Republic of Belarus)</i>	34
Karpenka A.F. <i>On the state of radioactively contaminated territories and their revival</i>	38
Kontsevaya I.I. <i>Influence of beta-lactam antibiotics on organogenesis in tissue culture of Karelian birch leaf explants</i>	43
Lazareva M.S., Klimovich L.K., Klimov A.V. <i>Features of the formation of oxalis oak forest</i> ..	50
Loginov D.N. <i>Species structure of the imago of blood-sucking mosquitoes (Diptera, Culicidae) in the Brest and Gomel regions of the Republic of Belarus</i>	56
Makarenko T.V. <i>Influence of natural sorbents on the accumulation of heavy metals in bottom sediments of water bodies</i>	62
Ostrovskij A.M. <i>Determinant of synanthropic woodlice species (Isopoda: Oniscidea) of the south-east of Belarus</i>	69
Sakovich V.V., Zhernossekov D.D. <i>Milk-clotting enzymes of various origin: prospects for application in cheese making</i>	75
Sokolov A.S. <i>Assessment of the landscape representativeness of the system of specially protected natural areas of the Belarusian Upland Landscape Province</i>	81
Uglyanec A.V., Garbaruk D.K. <i>Contamination by ¹³⁷Cs and ⁹⁰Sr of the dominant species of the living ground cover in the black alder forests of the exclusion zone of the Chernobyl NPP</i> ..	85
Khramchankova V.M., Matveyenkau M.V. <i>Cytotoxic and photomodifying properties of methanol and acetone extracts of five lichen species</i>	92

INFORMATION TECHNOLOGY

Voruev A.V., Demidenko I.O., Kolaib S.M., Domaskanov D.V. <i>Building a workplace for testing virtual network infrastructure</i>	99
Osipenko N.B., Mironenko V.M. <i>Automation of the selection of personalized sentences on the example of the problem of finding a movie</i>	105

MATHEMATICS

Borodich R.V., Selkin M.V. <i>On intersections of non-\mathfrak{F}-subgroups separated by a subgroup functor</i>	111
Galmak A.M., Selkin M.V. <i>Skew elements in polyadic groups of a special kind of odd arity. II</i>	118
Malinkovskij Yu.V. <i>Ergodicity criterion for one class of Markov chains with continuous time</i>	125
Mozhej N.P. <i>Equiaffine connections of zero curvature on homogeneous spaces with a solvable transformation group</i>	131
Nemilostivaya V.A. <i>Stationary distribution of networks with positive and negative claims and two types of nodes</i>	138
Pilipchuk L.A., Pilipchuk A.S., Polyachok E.N. <i>Two-criteria stream programming problems</i> .	144

PHYSICS

Leonenko D.V. <i>Transverse bending of a circular sandwich plate of stepped thickness</i>	151
Ohrimenko I.P., Kolchevskij N.N., Petrov P.V. <i>Model of a geostationary gravitational wave detector</i>	156

ABSTRACTS

Savarin A.A. <i>On the finding of the small white-toothed shrew (<i>Crocidura suaveolens</i>) on the territory of the Mogilev region of Belarus</i>	163
---	-----

UDC 582.287.238:57.033

Milk-clotting enzymes of various origin: prospects for application in cheese making

V.V. SAKOVICH, D.D. ZHERNOSSEKOV

The results of the comparative characteristics of rennet preparations have been presented. As the object of the study enzyme preparations possessing milk-clotting activity that are available in the Republic of Belarus have been considered. Indicators having important influence on the formation of a cheese curd were determined.

Keywords: rennet, cheese curd, physical and chemical properties.

В работе представлены результаты сравнительной характеристики ферментных препаратов сычужного действия. В качестве объекта исследования рассматривались доступные в Республике Беларусь ферментные препараты, обладающие молокосвертывающей активностью. Определялись показатели, оказывающие важное влияние на образование сырного сгустка.

Ключевые слова: сычужный фермент, сырный сгусток, физико-химические свойства.

According to the main Department of the processing industry of the Ministry of Agriculture and Food of the Republic of Belarus, the cheese market in the Republic of Belarus is constantly growing and developing branch. New types of cheese are being mastered; capacities for their production are increasing. The volume of production is 5 times the volume of consumption. In 2019 the Republic of Belarus exported cheese to 19 countries. In the structure of milk and dairy products supplies to the external market, cheeses account for almost 40 % of export earnings. The growth rate of cheese exports in 2019 was 118 %. The success of Belarusian dairy companies in the market is a combination of two factors: continuous work to improve the quality of products and affordable price for consumers [1].

According to the Republican unitary enterprise (RUE) «National Center for Marketing and Price Study» the main part of cheese is produced with the rennet. These enzymes are supplied by the Moscow rennet plant. There are also some European milk-clotting enzymes (from Denmark, Holland, Italy and other countries) which are supplied to Belarus by various intermediary firms [2]. Therefore, it is important to produce a domestic milk-clotting enzyme that will reduce the cost of purchasing rennet abroad.

The most famous enzyme traditionally used for milk coagulation is the rennet rennin or chymosin (EC 3.4.23.4). The enzyme belongs to the class of aspartate proteinases [3, p. 156], it is found in the juice of the fourth stomach of calves [4, p. 50], [5 p. 153].

The rapid expansion of dairy production and the shortage of animal rennet contributed to the emergence of new coagulants for milk of microbial and plant origin. Preparations of milk clotting enzymes can be conditionally divided into two groups: first – rennet; the second is rennet substitutes. The second group includes substitutes of animal, plant, microbial origin and recombinant chymosin.

Milk clotting enzyme preparations must meet the following basic criteria:

- strict compliance with the declared clotting activity of milk to the norm and, as a result, optimal consumption of preparations;
- high stability during long-term storage;
- optimal ratio of proteolytic and milk-clotting activity – one of the main characteristics that affect the quality of cheese;
- minimal non-specific proteolytic activity, which allows avoiding protein losses during cheese production, as well as the appearance of unpleasant tastes and consistency defects during cheese ripening.

Optimal thermal stability, allowing to strictly adhering to the main technological parameters of cheese production; – high purity of preparations [6 p. 50].

Not all milk-clotting enzymes meet these requirements and are therefore unsuitable for cheese making.

All commercial milk clotting enzymes are aspartate proteases (EC 3.4.23.) They specifically cleave the Phe105 – Met106 bond of bovine kappa casein.

The steady growth in global cheese production keeps the search for suitable rennet substitutes as the actual problem of science.

Fungi are an attractive source of various biologically active compounds [7, p. 1020]. Fungal enzymes are often produced in submerged conditions, but solid phase fermentation has also been used successfully. The cultivation conditions and the composition of the nutrient media strongly influence the ratio of milk clotting activity to the proteolytic activity of the enzyme [8, p. 997].

For each type of cheese, the use of one of the enzyme preparations offered by manufacturers should be theoretically and experimentally justified. Since in the Belarusian market there is a range of milk – clotting preparations that have different technological characteristics, we have attempted to give a comparative characteristic of rennet proteases presented on the Belarusian market of enzyme preparations.

Materials and methods. The following enzyme preparations were used in the work: rennet calf enzyme Renco; natural lamb enzyme Rennet; calf rennet E160; microbial fluid enzyme May-san; microbial enzyme Rennet; enzyme preparation from the culture fluid of *Pleurotus ostreatus*.

Protein concentration was determined spectrophotometrically [9 p. 356].

Milk-clotting activity was determined according to the modified Pyatnitsky method. The unit of milk-clotting activity (MCA) was defined as the amount of enzyme that clots 100 ml of milk in 40 min at 35 °C [10 p. 80].

The total proteolytic activity (PA) was determined by the lysis of gelatin in a thin layer of agar gel [11 p. 103]. For 1 unit of activity (E) of the enzyme was taken such an activity that leads to hydrolysis of the substrate in the area of the gel size 1 cm².

pH optimum of enzymes possessing milk clotting activity was measured at different pH values (from 3,5 to 7,5), using 0,2 M acetate buffer and 0,2 M phosphate buffer.

Temperature optimum of the activity of milk-clotting enzymes was determined after incubating the reaction mixture at different temperatures in the range from 25 to 45°C

The total proteolytic activity (PA) was determined by the lysis of gelatin in a thin layer of agar gel. For 1 unit of activity (E) of the enzyme was taken such an activity that leads to hydrolysis of the substrate over a gel area of 1 cm².

The pH optimum of enzymes with milk clotting activity was measured at various pH values (from 3,5 to 7,5) using 0,2 M acetate buffer and 0,2 M phosphate buffer.

The temperature optimum for the activity of milk coagulation enzymes was determined after incubation of the reaction mixture at various temperatures in the range from 25 to 45°C.

The effect of CaCl₂ concentration on MCA: enzyme activity was measured adding in milk various concentrations of CaCl₂ (from 2,5 to 30 mM).

The cheese curd formation was carried out in accordance with GOST 92225.

Results and discussion. The aim of the work was to study the activity of milk-clotting enzymes of different origin, which are used for the cheese production. The following preparations were chosen:

1. Rennet calf enzyme Renco. Natural rennet calf enzyme, a liquid salt solution comprising at least 92 % chymosin as the active component. Made in New Zealand («Renco»).

2. Natural lamb enzyme Rennet. This product is used for the production of fresh and medium-ripening cheese with a sweet and spicy taste. It is also very suitable for hard cheeses with a long ripening period. Made in Italy (Caglio Bellucci s.l.).

3. Calf rennet E160. (In the form of solution). The preparation is suitable for the production of any type of cheese – hard cheeses, semi-hard cheeses, soft and blue cheeses, as well as cottage cheese. Composition: Chymosin 80 %, Pepsin 20 %. Made in Italy (Caglio Bellucci s.l.).

4. Microbial enzyme Rennet. Milk-clotting enzyme preparation is the microbial coagulant and is produced on the basis of *Rhizomucor miehei*. All ingredients used in every stage of the manufacturing process are not animal-based. It is used for the production of fresh and medium-ripening cheeses. Enzyme composition: *chymosin*: 75 % – *pepsin*: 25 %. Made in Italy ((Caglio Bellucci s.l.).

5. Liquid microbial enzyme Mayasan. This food microbial enzyme is produced by controlled fermentation of *Rhizomucor miehei*. It is used for the preparation of soft cheeses, feta cheese, as well as semi-hard and hard cheese with a ripening period of 3–6 months. Made in Turkey (Istanbul, MAYASAN GIDA SAN).

6. An enzyme preparation developed at the Department of Biotechnology of the EE «Polesky State University» on the basis of the culture fluid of *Pleurotus ostreatus*.

Each preparation has the declared clotting activity, these data were compared with the results determined experimentally, at standard conditions according to GOST10 288-2001 «Preparations of milk clotting enzymes. Technical conditions». The results of these studies are presented in table 1.

Table 1 – Milk-clotting activity of commercial enzyme preparations

Trade mark	Milk-clotting activity, arbitrary units	
	The declared	Installed
Renco	580	580
Rennet	280	260
Rennet rhizomucor	700	750
E160	150	280
Mayasan	10000	1000

However, the claimed MCA data do not always coincide with those obtained experimentally. The experimental data results are presented in table 2.

Table 2 – MCA data of the studied enzyme preparations

Enzyme preparation	Volume, ml	Protein, mg/ml	Activity, units		Specific activity, unit/mg protein		MCA/PA
			MCA	PA	MCA	PA	
Renco	10	10	580	497	58	49,7	1,17
Rennet	10	10	260	265	26	26,5	0,98
Rennet rhizomucor	10	10	750	430	75	43	1,7
E160	10	10	280	191	28	19,1	1,5
Mayasan	10	10	1000	418	100	41,8	2,4
Enzyme preparation from <i>P. ostreatus</i>	1	1,07	81,08	6,32	75,78	5,91	13,7

The highest specific MCA (100 units/mg protein) contains preparation Mayasan, however, this preparation has high proteolytic activity, which is undesirable for rennet. It is known that with a high proteolytic activity of an enzyme preparation, not only clot formation is observed, but also its further hydrolysis. This leads to the appearance of bitter peptides and makes such an enzyme preparation not very suitable for use in cheese making [12, p. 49].

The enzyme preparations from *P. ostreatus* and Rennet Rhizomucor have almost identical specific MCA (about 75 U / mg protein), however, the PA index in the case of Rennet Rhizomucor is significantly higher than that of the enzyme preparation from *P. ostreatus*.

The lowest MCA values are shown for Rennet and E160 preparations (26 and 28 U/ mg protein correspondingly), which can negatively affect the cheese quality and significantly increase the time for its formation.

Influence of temperature. In order to study the physicochemical properties of milk-clotting preparations, the effect of temperature on the MCA was investigated. The effect of temperature on the catalytic activity of enzymes possessing MCA showed a typical activity-temperature dependence of the studied enzymes. The increased milk clotting activity at a higher temperature can be explained by the aggregation of proteins and molecular rearrangements in the protein structure [13 p. 345]. As it is shown in figure 2, for all enzyme preparations MCA was observed in the entire investigated temperature range from 25 to 45⁰ C. At the same time, the temperature optimum of the preparations is different.




Figure 1 – Effect of temperature on enzyme preparations

It was shown the different temperature optima for the studied enzymes, mainly because of their structural peculiarities. Conformational changes in the protein structure at high temperatures can make it vulnerable to proteolysis, since unfolding of the protein can present new cleavage sites to enzymatic hydrolysis [14, p. 395], [15, p. 483], [16, p. 1902]. For enzyme preparations from Rennet Rhizomucor, *P. ostreatus*, E160 and Renco, the temperature optimum is at 35⁰ C. This value is slightly lower than for the enzymes from Maysan and lamb Rennet. This is in accordance with literature data concerning temperature influence on the activity of enzyme preparations possessing MCA [17, p. 597].

Effect of pH. All enzyme preparations retained MCA in the pH range from 4 to 9. The pH optima of enzyme preparations from different origins are somewhat different (see figure 2).




Figure 2 – pH optima of enzyme preparations from different origins

According to Demir [18, p. 286], enzymes with high pH optima are more profitable in food production, because the forming clots are bad and weak at a lower pH of milk. The range of stability of milk-clotting proteinases is in the pH range from 3,5 to 7,5 [19, p. 527]. These data are in accordance with our results.

The influence of calcium. There are numerous data that calcium at a sufficiently high concentration is an important component in the formation of milk clot. The hardness of the cheese can be increased up to 81 % by the addition of about 10 mM CaCl₂, but higher concentrations of calcium chloride may decrease the hardness of the final product [20, p. 1393]. The use of high concentrations of calcium chloride can have negative consequences in cheese production. High calcium concentrations change the cheese fusibility, which causes a number of problems in cheese production. As it is shown in figure 3, the maximum MCA of the enzyme from *P. ostreatus* in our study was the highest when calcium chloride was added to the substrate (milk) at a final concentration of 10 mM, which is consistent with the literature data.




Figure 3 – Effect of calcium chloride concentration on enzyme preparations

For the Rennet Rhizomucor, E160, Rennet and Renco preparations, the highest MCA was obtained by adding 25 mM calcium chloride in final concentration that may have negative effect on cheese curd formation.

Cheese curd. According to GOST 92225, the quality of milk or enzyme suitability for cheese production are estimated by the nature of the formed cheese curd. The data of cheese curd quality with the studied enzyme preparations are presented in table 3.

Table 3 – Quality of the cheese curd

Enzyme preparation	Clot characteristic	Serum characteristics
Renco	The clot has a delicate, uniform consistency that meets the requirements for soft cheeses.	The serum does not have a bitter taste, it is transparent.
Rennet	The torn clot is soft to the touch with breaks. This clot is of satisfactory quality.	The serum is turbid.
Rennet rhizomucor	The clot has a delicate, uniform consistency that meets the requirements for soft cheeses.	The serum does not have a bitter taste, it is transparent.
E160	The torn clot is soft to the touch with breaks. This clot is of satisfactory quality.	The serum is turbid.
Mayasan	The clot is elastic to the touch, without eyes in the longitudinal section	The serum has a slightly bitter taste
Enzyme preparation from <i>P. ostreatus</i>	The clot has a creamy hue, uniform throughout the mass. The clot was elastic to the touch, without eyes in the longitudinal section. It has a light mushroom aroma.	The serum does not have a bitter taste, it is transparent.

Thus, the best cheese curd was obtained using the enzyme preparations of Renco, Rennet Rhizomucor, Mayasan and the enzyme preparation *P. ostreatus*. However, when using the Mayasan preparation, on serum had a slightly bitter taste, which was associated with the high proteolytic activity of the preparation.

Conclusion. This study was carried out to assess the competitiveness of milk-clotting enzymes of different origins. We have previously developed a scheme for the isolation and purification of the milk clotting enzyme from the culture liquid of *Pleurotus ostreatus*. This preparation has similar characteristics with these ones which are available in the market of Republic of Belarus: high milk clotting activity, the similar value of pH optimum (5,0) and temperature optimum (35°C), similar effect of 10 mM calcium chloride. The enzyme from *Pleurotus ostreatus* forms the cheese curd of high quality. This preparation also has its own peculiarities: a high ratio of milk-clotting and proteolytic activity, which is a good indicator for the formation of a cheese curd. Also, the cheese curd formed with the enzyme from *Pleurotus ostreatus* has a slight mushroom aroma.

Thus, it is possible to propose the enzyme preparation from the culture liquid of *Pleurotus ostreatus* for use in the dairy industry at the stage of cheese curd formation.

Literature

1. ProductBY [Electronic resource] : Belarusian food trade and industrial portal. – Mode of access : <https://produkt.by/news/mariya-klimova-belorusskoe-syrodellie-vchera-segodnya-zavtra>. – Date of access : 20.10.2020.
2. Infobase.BY [Electronic resource] : Transport, agriculture, industry in Belarus. – Mode of access : <https://www.infobaza.by/article/agro/obzorsyrov/>. – Date of access : 20.10.2020.
3. Starovoitova, V. V. Study of functional properties of calf chymosin and its recombinant forms : dis. ... cand. of chem. sciences : 02.00.10 / V. V. Starovoitova. – Moscow, 2001. – 157 p.
4. Kolesnikova, S. S. Enzymes for milk coagulation in cheese making / S. S. Kolesnikova // Dairy business. – 2006. – № 8. – P. 50–51.
5. Ayhan, F. The effect of fermentation parameters on the production of *Mucor miehei* acid protease in a chemically defined medium / F. Ayhan, S. S. Celebi, A. Tanyolac // J. Chem. Technol. Biotechnol. – 2001. – № 76. – P. 153–160.
6. Gorina, T. A. Innovations in the field of milk-converting enzymes / T. A. Gorina // Cheese and butter making. – 2009. – № 3. – P. 50–51.
7. Chopra, A. K. Factors affecting protease production by *Bacillus stearothermophilus* RM-67 / A. K. Chopra, D. K. Mathur // J. Food Protect. – 1983. – V. 46. – P. 1020–1025.
8. Vairo Cavalli, S. Hydrolysis of caprine and ovine milk proteins, brought about by aspartic peptidases from *Silybum marianum* flowers / S. Vairo Cavalli, S. V. Silva, C. Cimino, F. X. Malcata, N. Priolo // Food Chemistry. – 2008. – V. 106. – P. 997–1003.
9. Kim, J.H. Characterization of a metalloenzyme from a wild mushroom, *Tricholoma saponaceum* / J. H. Kim, Y. S. Kim // Biosci. Biotechnol. Biochem. – 2001. – V. 65 (2). – P. 356–62.
10. Pyatnitsky, N. P. Determination of the activity of chymotrypsin at a speed of milk has been curdled and strained-acetate mixtures / N. P. Pyatnitsky, M. T. Proskuryakov // 17th century. Conf. physiologists of the South of Russia. – Stavropol, 1969. – T. 2. – P. 80.
11. Leighton, T. J. The relationship of serine protease activity to RNA polymerase modification and sporulation in *Bacillus subtilis* / T. J. Leighton, R. H. Doi, R. A. J. Warren, R.A. Kelln // J. Mol. Biol. – 1973. – V. 76. – P. 103–122.
12. Neveen M. Purification and characterization of Milk -Clotting Enzyme from the edible mushroom *Pleurotus albidus* / M. Neveen, N. Mohamed [et al.] // Research Journal of Pharmaceutical, Biological and Chemical Sciences. – 2018. – V. 9(5). – P. 49.
13. Najera, A. I. Effects of pH, temperature, CaCl₂ and enzyme concentrations on the rennetclotting properties of milk: a multifactorial study / A. I. Najera, M. De renobales, L. J. R. Barron // Food Chemistry. – 2003. – V. 80. – P. 345–352.
14. Mohamed Ahmed, I. A. Characterization of partially purified milk-clotting enzyme from *Solanum dubium* Fresen seeds / I. A. Mohamed Ahmed, I. Morishima, E. E. Babiker, N. Mori // Food Chemistry. – 2009. – V. 116 (2). – P. 395–400.
15. Mohamed Ahmed, I. A. Dubiumin, a chymotrypsin-like serine protease from the seeds of *Solanum dubium* Fresen / I. A. Mohamed Ahmed, I. Morishima, E. E. Babiker, N. Mori // Phytochemistry. – 2009. – V. 70 (4). – P. 483–491.
16. Mazorra-Manzano, M. A. Comparison of the milk-clotting properties of three plant extracts / M. A. Mazorra-Manzano, T. C. Perea-Gutiérrez [et al.] // Food Chemistry. – 2013. – V. 141. – P. 1902–1907.
17. Rao, M. B. Molecular and biotechnological aspects of microbial proteases / M. B. Rao, A. M. Tanksale, M. S. Ghatge, V. V. Despande // Microbiology and Molecular Biology Reviews. – 1998. – V. 62. – P. 597–635.
18. Demir, Y. Cysteine protease (capparin) from capsules of caper (*Capparis spinosa*) / Y. Demir, A. Güngör, E. D. Duran, N. Demir // Food Technol. Biotechnol. – 2008. – V. 46. – P. 286–291.
19. Preetha, S. Influence of culture conditions on the production of milk-clotting enzyme from *Rhizomucor* / S. Preetha, R. Boopathy // World Journal of Microbiology & Biotechnology. – 1994. – V. 10. – P. 527–530.
20. Chazarra, S. Characterization of the milk-clotting properties of extracts from artichoke (*Cynara scolymus*, L) flowers / S. Chazarra, L. Sidrach, D. López-Molina, J. N. Rodríguez-López // Int Dairy J. – 2007. – V. 17. – P. 1393–1400.